

Magnetic-field-induced superconductivity and superfluidity of W and Z bosons: in tandem transport and kaleidoscopic vortex states

M. N. Chernodub,^{1,2,*} Jos Van Doorselaere,² and Henri Verschelde²

¹*CNRS, Laboratoire de Mathématiques et Physique Théorique, Université François-Rabelais Tours, Fédération Denis Poisson, Parc de Grandmont, 37200 Tours, France*

²*Department of Physics and Astronomy, University of Gent, Krijgslaan 281, S9, B-9000 Gent, Belgium*
(Dated: March 27, 2012)

We show that in a background of a sufficiently strong magnetic field the electroweak sector of the quantum vacuum exhibits superconducting and, simultaneously, superfluid properties due to magnetic-field-induced condensation of, respectively, W and Z bosons. The phase transition to the “tandem” superconductor–superfluid phase – which is weakly sensitive to the Higgs sector of the standard model – occurs at the critical magnetic field of 10^{20} Tesla. The superconductor–superfluid phase has anisotropic transport properties as both charged and neutral superflows may propagate only along the magnetic field axis. The ground state possesses an unusual “kaleidoscopic” structure made of a hexagonal lattice of superfluid vortices superimposed on a triangular lattice of superconductor vortices. A weak electric field will induce both superconducting and, unexpectedly, superfluid flows.

PACS numbers: 12.15.-y, 13.40.-f, 74.90.+n

It is known that an extremely high magnetic field of hadronic scale may lead to plenty of unusual effects both in (dense) matter and in the quantum vacuum. The chiral magnetic effect [1] provides a particularly interesting example: a charge-parity-odd matter may generate an electric current along the axis of the magnetic field [3]. The corresponding conditions may be realized in noncentral heavy-ion collisions [2] in which a hot quark matter is created in a background of extremely high magnetic fields [4, 5]. Similar conditions may have existed in the very early moments of our Universe [9]. The strong magnetic field affects phases of the cold dense matter in the cores of strongly magnetized neutron stars [6].

Because of quantum effects an empty space may also exhibit quite unusual properties in a sufficiently strong magnetic background. In the background of a relatively low magnetic field of QED scale the vacuum should become optically birefringent [7]. The hadron-scale magnetic field should lead to magnetic catalysis [8], which implies, in particular, a steady enhancement of the chiral symmetry breaking in QCD vacuum as the external magnetic field strengthens.

More recently it was found that the vacuum becomes an *electromagnetic* superconductor in sufficiently strong external magnetic fields [10, 11]. The superconductivity of, basically, empty space, is mediated via spontaneous creation of a (charged) ρ -meson condensate if the magnetic field exceeds the critical value of

$$B_c^{\text{QCD}} = M_\rho^2/e \simeq 1.0 \times 10^{16} \text{ Tesla}, \quad (1)$$

where $M_\rho = 775.5 \text{ MeV}$ is the mass of the ρ meson. The ground state of the vacuum superconductor is characterized by inhomogeneous ground state of a very particular

geometric structure [12], possessing intriguing metamaterial (“perfect lens”) properties [13]. The magnetic fields of the required strength (1) may be created on Earth in heavy-ion collisions at the Large Hadron Collider [5].

We show that as the background magnetic field strengthens further, the Standard Model experiences a second superconducting, and, simultaneously, superfluid transition associated with a condensation of the W and Z bosons at a larger critical magnetic field:

$$B_c^{\text{EW}} = M_W^2/e \simeq 1.1 \times 10^{20} \text{ Tesla}, \quad (2)$$

where $M_W = 80.4 \text{ GeV}$ is the mass of the W boson. The condensation of the W bosons at the magnetic field higher than the critical value (2) was predicted by Ambjørn and Olesen in Refs. [14]. The key idea here is that the vacuum of charged vector particles (i.e., of the W mesons) is unstable in the background of a sufficiently strong magnetic field provided these particles have anomalously large gyromagnetic ratio $g_m = 2$. The large value of g guarantees that the magnetic moment of such particles is too large to withstand a spontaneous condensation at sufficiently strong external magnetic fields. We show that the W condensate leads to exotic superconducting and superfluid effects.

The bosonic part of the electroweak sector of the Standard Model is described by the Lagrangian,

$$\mathcal{L} = -\frac{1}{4}W_{\mu\nu}^a W^{a,\mu\nu} - \frac{1}{4}X_{\mu\nu} X^{\mu\nu} + (D_\mu \Phi)^\dagger (D^\mu \Phi) - \lambda (|\Phi|^2 - v^2/2)^2. \quad (3)$$

where Φ is the complex Higgs doublet which interacts with $SU(2)_L$ and $U(1)_X$ gauge fields (W_μ^a and X_μ , respectively) via the covariant derivative $D_\mu = \partial_\mu - ig\tau^a W_\mu^a/2 - ig'X_\mu/2$, and τ^a are the Pauli matrices. The corresponding field strengths are $W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g\epsilon^{abc}W_\mu^b W_\nu^c$ and $X_{\mu\nu} = \partial_\mu X_\nu - \partial_\nu X_\mu$.

* On leave from ITEP, Moscow, Russia.




FIG. 1. (a) The W condensate $\langle |W|^2 \rangle^{1/2}$; (b) the gain in the energy density (6) due to the condensation, $\delta E = \langle E \rangle_W - \langle E \rangle_{W=0}$, (c) the Higgs expectation value $\langle \Phi^\dagger \Phi \rangle^{1/2}$. Eq. (17), and (d) the superconductivity transport coefficient (20) vs. the magnetic field B at fixed values of the Higgs masses $M_H = (1.1 \dots 2.0)M_Z$ (all quantities are averaged over the transversal plane).

The mexican hat potential in Eq. (3) breaks the electroweak symmetry down to the electromagnetic subgroup, $SU(2)_L \times U(1)_X \rightarrow U(1)_{\text{em}}$ because the Higgs field Φ acquires a quantum expectation value, $\langle \Phi \rangle \neq 0$. In the unitary gauge, $\langle \Phi \rangle = (0, v)^T$, the third component of the non-Abelian gauge field W_μ^3 mixes with the Abelian gauge field X_μ providing us with the massive Z_μ boson and the massless electromagnetic field A_μ : $W_\mu^3 = \sin \theta A_\mu + \cos \theta Z_\mu$, $X_\mu = \cos \theta A_\mu - \sin \theta Z_\mu$ where θ is the electroweak mixing (Weinberg) angle with $e = g \sin \theta = g' \cos \theta$ being the electric charge.

The classical equations of motion are:

$$\begin{aligned} 0 &= \partial^\mu W_{\mu\nu}^a + g\epsilon^{abc}W_\mu^b W_\nu^c - ig[(D_\mu \Phi)^\dagger \tau^a \Phi - h.c.]/2, \\ 0 &= \partial^\mu Y_{\mu\nu} - ig'(D_\mu \Phi)^\dagger \Phi - h.c./2, \\ 0 &= -D_\mu D^\mu \Phi + 2\lambda \Phi(|\Phi|^2 - v^2/2), \end{aligned} \quad (4)$$

The instability of the vacuum in the presence of the sufficiently strong magnetic field was clearly demonstrated by Ambjørn and Olesen in Ref. [14], and we briefly repeat their arguments here. Consider a uniform time-independent magnetic field directed along the third axis, $B_{\text{ext},i} = B_{\text{ext}}\delta_{i3}$ (we always take $eB > 0$ for simplicity of our notations). Then the quadratic part of the transverse (with respect to the magnetic field axis) components of the $W_\mu \equiv W_\mu^-$ field in Eq. (3) reads

$$\delta \mathcal{L}_{W_\perp}^{(2)} = (W_1^\dagger, W_2^\dagger) \begin{pmatrix} M_W^2 & -ieB_{\text{ext}} \\ ieB_{\text{ext}} & M_W^2 \end{pmatrix} \begin{pmatrix} W_1 \\ W_2 \end{pmatrix}, \quad (5)$$

(with $M_W = gv/2$), while the mass terms of the longitudinal components W_3 and W_0 , and of other vector particles are not affected at the classical level. The mass eigenvalues of Eq. (5) are $\mu_\pm^2 = M_W^2 \pm eB$. One of the masses, μ_- , vanishes at the critical value B_c of the magnetic field (2). This mass becomes purely imaginary at $B > B_c$, thus signaling a tachyonic instability towards condensation of the transverse components of the W_μ field, with the unstable eigenvector $(W_1, W_2) = (W, -iW)/2$, where W is a scalar field.

Since we consider the solutions in the transverse (x_1, x_2) plane for the transverse components of the fields, it is very convenient to use the complex notations for the coordinates, $z = x_1 + ix_2$, and for the vectors $\mathcal{O}_\mu = \partial_\mu, A_\mu, Z_\mu, W_\mu$ and their field strengths: $\mathcal{O} =$

$\mathcal{O}_1 + i\mathcal{O}_2$, $\bar{\mathcal{O}} = \mathcal{O}_1 - i\mathcal{O}_2$ and $\mathcal{O}_{12} = -\frac{i}{2}(\bar{\mathcal{O}}\mathcal{O} - \partial\bar{\mathcal{O}})$. Notice that $W^\dagger \neq \bar{W}$.

We use a symmetric gauge for the external magnetic field, $A_{\text{ext},1} = -Bx_2/2$ and $A_{\text{ext},2} = Bx_1/2$, so that the corresponding covariant derivative is $\mathcal{D}_{\text{ext}} \equiv \partial - ieA_{\text{ext}} = \partial + ezB_{\text{ext}}/2$. The $\bar{W}$ component of the W^- corresponds to the μ_+ eigenvalue of the operator in Eq. (5), so that it is not condensed. Thus, we put $\bar{W} = 0$ as it does not lower the energy, and continue to work in the unitary gauge $\Phi = (0, \phi)^T$, where ϕ is a real-valued field.

In complex notations the energy density is:

$$\begin{aligned} E &= \frac{1}{2} |(\bar{\mathcal{D}} + ig \cos \theta \bar{Z})W|^2 + \frac{1}{2} Z_{12}^2 + \frac{1}{2} B^2 + \frac{g^2}{8} |W|^4 \\ &+ \frac{1}{2} (-eB - g \cos \theta Z_{12} + \frac{g^2}{2} \phi^2) |W|^2 + \frac{g^2}{4 \cos^2 \theta} |Z|^2 \phi^2 \\ &+ \bar{\partial} \phi \partial \phi + \lambda (\phi - v^2/2)^2, \end{aligned} \quad (6)$$

and the equations of motion then become:

$$\mathfrak{D} \bar{\mathfrak{D}} W = \left(\frac{g^2}{2} |W|^2 - g \cos \theta Z_{12} - eF_{12} + \frac{g^2}{2} \phi^2 \right) W, \quad (7)$$

$$\bar{\mathfrak{D}}^2 W = 0, \quad (8)$$

$$\bar{\partial} F_{12} = \frac{e}{2} \bar{\partial} |W|^2 + \frac{e}{2} W^\dagger \bar{\mathfrak{D}} W, \quad (9)$$

$$0 = \cos \theta \bar{\partial} F_{12} - \sin \theta \bar{\partial} Z_{12} + ig^2 \frac{\sin \theta}{2 \cos^2 \theta} \bar{Z} \phi^2 \quad (10)$$

$$\partial \bar{\partial} \phi = \frac{g^2}{4 \cos^2 \theta} |Z|^2 \phi + \frac{g^2}{4} |W|^2 \phi + 2\lambda \phi (\phi^2 - \frac{v^2}{2}), \quad (11)$$

$$0 = \phi \bar{\mathfrak{D}} W + 2W \left(\bar{\partial} + i \frac{g}{2 \cos \theta} \bar{Z} \right) \phi + c.c., \quad (12)$$

where $\mathfrak{D} = \mathcal{D} + ig \cos \theta Z$ is a covariant derivative.

In Ref. [14] the equations of motion are solved in the Bogomolny limit, $M_Z = M_H$, where $M_H = \sqrt{2}\lambda v$ and $M_Z = gv/(2 \cos \theta)$ are the masses of the Higgs and Z bosons, respectively. Here we solve – partially following Ref. [15] – the equations of motion for arbitrary mass of the Higgs in the region $B \geq B_c$ close to the phase transition point, $|B - B_c| \ll B_c$. The later condition implies that the quantity $\epsilon = |W|/M_W \ll 1$ can serve as a small expansion parameter.

The combination of the two equations of motion, (7) and (8), with the requirement of the minimization of the energy density (6), lead to the simple Abrikosov equation,

$$\bar{\mathfrak{D}} W \approx \bar{\mathfrak{D}} W = 0, \quad (13)$$




FIG. 2. (a) The superconducting W condensate (14); (b) the superfluid Z condensate (17); (c) the Higgs expectation value (17); and (d) the superconducting (20) and (e) superfluid (21) transport coefficients as the functions of the transverse plane coordinates x_1 and x_2 at the Higgs mass $M_H = 1.1M_Z$ (the solid surfaces) in the background magnetic field $B = 1.01B_c^{\text{EW}}$ directed along the x_3 axis. The semitransparent surfaces correspond to $M_H = 2M_Z$ in Figs. (a), (b), (c) and (d); and to $\kappa^Z = 0$ plane in Fig. (d). The red line in Fig. (c) gives the $B = 0$ Higgs expectation value, $\phi = v/\sqrt{2}$. Figure (f) illustrates the 3d regions of space predominantly occupied by the superconducting electric current J_3^E of the W bosons and the superfluid neutral flows J_3^Z of the Z bosons generated by a weak test electric field E_{ext} parallel to the strong magnetic field $B = 1.01B_c^{\text{EW}}$ at $M_H = 1.1M_Z$.

valid up to corrections of the order of $O(\epsilon^2)$. This equation has nontrivial periodic solutions known as Abrikosov lattices. Following Abrikosov [16], we choose a general solution of Eq. (13) as a sum over lowest Landau levels:

$$W(z) = \sum_{n \in \mathbb{Z}} C_n e^{-\frac{\pi}{2}(|z|^2 + \bar{z}^2) - \pi\nu^2 n^2 + 2\pi\nu n \bar{z}}, \quad (14)$$

where $L_B = \sqrt{2\pi/(eB)}$ is the magnetic length and ν is an arbitrary real valued parameter. In order to ensure a regular structure of the lattice, the complex coefficients C_n are usually chosen in a periodic manner: $C_{n+N} = C_n$.

The solution with $N = 1$ and $\nu = 1$ defines the square lattice of the original Abrikosov's solution [16]. However, the global energy minimum is reached for the equilateral triangular lattice at $N = 2$ with $C_1 = \pm iC_0$ and $\nu = \sqrt[4]{3}/\sqrt{2} \approx 0.9306$ in agreement with earlier studies devoted to the W condensation [14, 15]. The energy density (6) is then minimized numerically for fixed values of the magnetic field B and the Higgs mass M_H [the later is chosen in a phenomenologically interesting range of $M_H = (1.1 \dots 2.0)M_Z$]. At $B \geq B_c^{\text{EW}}$ the condensation of the W bosons, Fig. 1(a), makes the energy density smaller compared to its value in the trivial ground state, Fig. 1(b). The heavier the Higgs boson the weaker the effect of the magnetic field on condensates and, as we will see, on transport coefficients.

Equations (13) and (9) imply $\partial(B - e|W|^2/2) = 0$, up to $O(\epsilon^2)$ terms. The solution carrying a finite energy per

unit cell $\mathcal{A}$ of the Abrikosov lattice is

$$B(z) = B_{\text{ext}} + \frac{e}{2}|W(z)|^2 - \frac{e}{2} \frac{1}{\text{Area}(\mathcal{A})} \int_{\mathcal{A}} dz d\bar{z} |W|^2, \quad (15)$$

where the integration constant – given by the integral over the unit lattice cell $\mathcal{A}$ – guarantees the conservation of the magnetic flux, $\int_{\mathcal{A}} dz d\bar{z} B(z) = \text{Area}(\mathcal{A}) \cdot B_{\text{ext}}$. Thus, the magnetic field (15) becomes transversally non-uniform due to the backreaction of the W condensate.

Then, the solution of Eqs. (10) and (11), are given by the nonlocal expressions for the Z and Higgs condensates:

$$Z \equiv Z_1 + iZ_2 = -i \frac{g \cos \theta}{2} \frac{\partial_1 + i\partial_2}{-\Delta + M_Z^2} |W|^2, \quad (16)$$

$$\phi = \frac{v}{\sqrt{2}} \left(1 - \frac{g^2}{4} \frac{1}{-\Delta + M_H^2} |W|^2 \right) \quad (17)$$

respectively, with $\Delta \equiv \bar{\partial}\partial = \partial_1^2 + \partial_2^2$. The remaining equation (12) is satisfied automatically up to $O(\epsilon^2)$.

In the ground state the W , Z and Higgs condensates are functions of the transversal coordinates x_1 and x_2 , as visualized in Figs. 2(a), (b) and (c), respectively. The expectation value of the Higgs field falls down as the magnetic field rises, with a slope which becomes weaker as the Higgs mass rises, Figs. 1(c) and 2(c).

We point out that the ground state of the vacuum at $B > B_c^{\text{EW}}$ is a “tandem” phase which is, simultaneously, an electromagnetic superconductor and a neutral superfluid. Indeed, introducing an infinitesimally weak test




FIG. 3. Kaleidoscopic ground state: the density plots of the phases (left) of the W condensate (14) and (right) of the Z condensate (17) in the transversal (x_1, x_2) plane at $B = 1.01B_c^{\text{EW}}$. The endpoints of the cuts in the phases (shown by the circles) are the superconductor vortices and superfluid (anti)vortices, respectively.

electric field E^{ext} one can prove – with the use of the equations of motion (4) – the following transport laws for the electromagnetic current $J_\mu^E = \partial^\nu F_{\nu\mu} \propto \delta\mathcal{L}/\delta A^\mu$ and the neutral Z -boson current $J_\mu^Z = \partial^\nu Z_{\nu\mu} \propto \delta\mathcal{L}/\delta Z^\mu$, respectively:

$$\partial_{[0} J_{3]}^E(x) = -\kappa^E(x_1, x_2) \cdot E_3^{\text{ext}}, \quad \partial_{[0} J_{i]}^E = 0, \quad (18)$$

$$\partial_{[0} J_{3]}^Z(x) = -\kappa^Z(x_1, x_2) \cdot E_3^{\text{ext}}, \quad \partial_{[0} J_{i]}^Z = 0, \quad (19)$$

where $i = 1, 2$. The transport parameters for the electromagnetic and neutral currents are, respectively, as follows,

$$\kappa^E(x_1, x_2) = e^2 |W|^2(x_1, x_2), \quad (20)$$

$$\kappa^Z(x_1, x_2) = -e^2 \cot \theta \frac{\Delta}{-\Delta + M_Z^2} |W|^2(x_1, x_2). \quad (21)$$

They are functions of the transversal coordinates $x_{1,2}$, as shown in Figs. 2(d) and (d), respectively.

Equation (18) implies anisotropic superconductivity of the ground state at $B > B_c^{\text{EW}}$ similarly to an analogous phenomenon in QCD [10, 11]: a weak electric field introduces a resistance-free growth of electric current which continues streaming after the field is switched off. Equation (19) implies an anisotropic superfluidity of the neutral Z currents, and, simultaneously, it illustrates a very unusual physical effect: an external electric field induces a superfluid current of neutral particles which are flowing frictionlessly along the magnetic field axis.

From the point of view of the electric conductivity properties, a ground state of the vacuum can either be a superconductor or an insulator due to Lorentz symmetry (indeed, a dissipative behavior, like in the Ohm law, is inconsistent with the Lorentz symmetry of the vacuum). Thus, the absence of the electric resistance (and vanishing shear and bulk viscosities) in the $B > B_c^{\text{EW}}$ phase are protected by a remnant Lorentz symmetry in the (x_0, x_3) plane. Similar Lorentz-protection arguments apply to the superfluid property.

The superconductivity coefficient (20), averaged over the transversal (x_1, x_2) plane, is a linearly growing function of the magnetic field B , Fig. 1(d), at $B > B_c^{\text{EW}}$. The superfluid coefficient (21) is a sign-changing function, Fig. 2(e), of the transversal coordinates $x_{1,2}$ which has vanishing mean value if averaged over the transversal plane. Thus, an external electric field E_3^{ext} gives rise to (i) a nonzero net electric current along the magnetic field axis and (ii) a neutral superfluid space-dependent flow in opposite directions with vanishing net current, Fig. 2(f). The superconductivity and superfluidity always appear in a tandem. The transversal components $E_{1,2}^{\text{ext}}$ induce neither superconducting nor superfluid currents.

It is known that the ground state of the vacuum at $B > B_c^{\text{EW}}$ is an equilateral triangular lattice of the vortex defects in the W field [14, 15] (we call these vortices the “superconductor vortices”). At the vortex positions the field $W \propto W_1^- + iW_2^-$ vanishes, Fig. 2(a), and its phase, $\arg(W)$, winds around the vortex position. We find that the ground state has a much more complicated structure in the neutral sector: the state has a hexagonal lattice of the “superfluid” vortices (and antivortices), characterized by the vanishing $Z \equiv Z_1 + iZ_2$ field, Fig. 2(b), and positive (negative) winding numbers [17]. The combined “kaleidoscopic” pattern of the vortex lattices, superimposed on the density plots of the phases of the superconducting, W and the superfluid, Z , fields, are shown in Fig. 3. All superconductor vortices are always located on top of certain superfluid vortices.

In conclusion, in this paper we have unravelled the detailed structure of the electroweak sector of the vacuum in a strong magnetic field and discovered a kaleidoscopic pattern of superconducting W -boson vortices and superfluid Z -boson vortices and antivortices. We have shown that application of an electric field generates a superconducting current in tandem with neutral superfluid currents. The neutral currents flow predominantly along the Z -boson vortices and have vanishing net current.

- [1] K. Fukushima, D. E. Kharzeev and H. J. Warringa, Phys. Rev. D **78**, 074033 (2008).
- [2] D. Kharzeev, Phys. Lett. B **633**, 260 (2006).
- [3] A. Vilenkin, Phys. Rev. D **22**, 3080 (1980).
- [4] V. Skokov, A. Y. Illarionov and V. Toneev, Int. J. Mod. Phys. A **24**, 5925 (2009).

- [5] W. T. Deng and X. G. Huang, arXiv:1201.5108 [nucl-th].
- [6] E. J. Ferrer, V. de la Incera and C. Manuel, Phys. Rev. Lett. **95**, 152002 (2005); S. Fayazbakhsh and N. Sadooghi, Phys. Rev. D **82**, 045010 (2010); A. Rabhi, P. K. Panda and C. Providencia, Phys. Rev. C **84**, 035803 (2011); A. A. Isayev and J. Yang, Phys. Rev. C **84**,

- 065802 (2011).
- [7] S. L. Adler, *Annals Phys.* **67**, 599 (1971).
 - [8] K. G. Klimenko, *Z. Phys. C* **54**, 323 (1992); V. P. Gusynin, V. A. Miransky and I. A. Shovkovy, *Phys. Rev. Lett.* **73**, 3499 (1994); V. P. Gusynin, V. A. Miransky and I. A. Shovkovy, *Nucl. Phys. B* **462**, 249 (1996).
 - [9] D. Grasso, H.R. Rubinstein, *Phys. Rept.* **348**, 163 (2001).
 - [10] M. N. Chernodub, *Phys. Rev. D* **82**, 085011 (2010).
 - [11] M. N. Chernodub, *Phys. Rev. Lett.* **106**, 142003 (2011).
 - [12] M. N. Chernodub, J. Van Doorselaere and H. Verschelde, *Phys. Rev. D* **85**, 045002 (2012).
 - [13] I. I. Smolyaninov, *Phys. Rev. Lett.* **107**, 253903 (2011).
 - [14] J. Ambjorn and P. Olesen, *Nucl. Phys. B* **315**, 606 (1989); *Phys. Lett. B* **218**, 67 (1989); *Int. J. Mod. Phys. A* **5**, 4525 (1990).
 - [15] S. W. MacDowell and O. Tornkvist, *Phys. Rev. D* **45**, 3833 (1992); O. Tornkvist, hep-ph/9204235.
 - [16] A. A. Abrikosov, *Sov. Phys. JETP* **5**, 1174 (1957).
 - [17] The superconductor and superfluid vortices should be distinguished from the existing W^- and Z^- electroweak vortex solutions [18], including known vortex solutions which carry electric currents. [19].
 - [18] See, e.g., a review in A. Achucarro and T. Vachaspati, *Phys. Rept.* **327**, 347 (2000).
 - [19] M. S. Volkov, *Phys. Lett. B* **644**, 203 (2007); J. Garaud and M. S. Volkov, *Nucl. Phys. B* **826**, 174 (2010).